

**Abstract**—Data from an existing site preparation experiment in the Georgia Piedmont were subjected to a modeling approach to analyze effects of site preparation intensity on stand development of loblolly pine (*Pinus taeda* L.) 5 to 12 years since treatment. An average stand height model that incorporated indicator variables for treatment provided an accurate description of responses to site-preparation intensity, with steepness of the height trajectory increasing with site-preparation intensity. Stand basal area and volume varied similarly among treatments as found for height, with their development increasing with treatment intensity.

## INTRODUCTION

It has been widely accepted that development of stand height follows a certain pattern that can be described by the following model (Clutter and others 1983):

$$\ln(H)=a+b/AGE \quad (1)$$

where

H is stand height,

AGE is stand age, and

a and b are estimated parameters.

Similarly, growth of stand basal area (BA) and stand volume (V) follow a similar pattern as:

$$\ln(Y)=a+b/AGE \quad (2)$$

where

Y is stand basal area or stand volume, and

a, b, and AGE are as defined previously.

The growth pattern of stand height, basal area, and volume can vary with site quality, species, and silvicultural treatments, such as site preparation. For a given species and site, effects of treatments can be detected by incorporating their effects into this model:

$$\ln(Y)=a+b/AGE+cTRMT \quad (3)$$

where

TRMT represents treatment effects,

c is an estimated parameter, and

a, b, and AGE are as defined previously.

In this study, an indicator variable was specified for each treatment to detect effects of site preparation intensity on stand development.

## DATA

The data were from an existing site preparation study of loblolly pine (*Pinus taeda* L.) initiated in 1980 in the Georgia Piedmont. After clearcutting mature loblolly pine, the six site-preparation intensities were applied, ranging from absence of site preparation to combinations of mechanical, herbicide, and fertilizer treatments. The treatments are listed as follows in order of increasing intensity:

- (1) Clearcut only.
- (2) Chainsaw. All residual trees greater than 2.5 centimeters diameter at breast height (d.b.h.) were felled with a chainsaw in August 1981.
- (3) Shearing of residual trees with a KG blade mounted on a D7 tractor in September 1981 and chopping of woody debris with a single pass of a rotary-drum chopper in November 1981.
- (4) Shear, chop, and herbicide. Treatment 3 plus application of 0.5 cubic centimeters Velpar (TM) Gridball pellets (hexazinone at 10 percent active ingredient) in a 0.6-meter x 0.6-meter grid pattern at a rate of 2.8 kilograms per hectare in March 1982.
- (5) Shear, rootrake, burn, and disk. Residual trees were sheared and rootraked into windrows in September 1981 and burned in October 1981. The remaining debris and ash were scattered with a dozer blade and the plots were disked with an offset harrow to a depth of 15-20 centimeters in October 1981.
- (6) Shear, rootrake, burn, disk, fertilize, and herbicide. Treatment 5 plus a broadcast application of ammonium- nitrate fertilizer at 114 kilograms N per hectare and a 1.2-meter band application of Oust (TM) (sulfometuron) at 0.42 kilograms active ingredient per hectare in March and April 1983.

Each treatment was replicated five times in a randomized complete-block design. In January and February 1982, seedlings of loblolly pine were hand-planted at a spacing of 1.8 meters x 3 meters. After establishment of the stands, measurements of d.b.h. (centimeters) and height (meters)

<sup>1</sup> Graduate Research Assistant and Assistant Professor, respectively, Daniel B. Warnell School of Forest Resources, University of Georgia, Athens, GA 30602-2152; and Research Ecologist, USDA Forest Service, Southern Research Station, Juliette, GA 31046 (respectively).

of each planted pine were taken 5, 8, 10, and 12 growing seasons after treatment, from which stand average height (H), basal area (BA) (square meters per hectare), and volume (V) (cubic meters per hectare) were calculated. All models were fitted with linear regression using a 95 percent significance level.

## RESULTS AND DISCUSSION

### Stand Average Height

Fitting stand average height to model (3) resulted in the following equation:

$$\ln(HT) = 3.26 - 9.84/AGE - 0.415T1 - 0.256T2 - 0.126T3 - 0.316T4 - 0.0991T5 \quad (4)$$

where

T1, T2, T3, T4, and T5 are indicator variables that represent treatments 1 to 5, respectively.

For example, T1=1 if treatment is 1, otherwise T1=0. Equation (4) indicates that development of average height differed significantly among treatments, with rate increasing with site-preparation intensity. Average height of untreated stands was significantly less than that of treated stands ( $P \leq 0.05$ ).

Numerous studies have reported increases in stand height in response to site preparation (Glover and Zutter 1993, Harrington and Edwards 1996, Pienaar and Rheney 1995, Thomson and McMinn 1989). Since stand height is relatively similar for a wide range of stand densities, increases in the rate of height development probably are more attributable to improvement in site quality due to site preparation. Measurements of soil properties on this site demonstrated that the treatments improved growing conditions for pine by decreasing bulk density and increasing pore space (Miller and Edwards 1985).

### Stand Basal Area

The following equation resulted from fitting stand basal area to model [3]:

$$\ln(BA) = 4.75 - 919/AGE - 1.52T1 - 1.04T2 - 0.308T3 - 0.477T4 \quad (5)$$

The fitted equation (5) indicates that development of stand basal area also increased with treatment intensity, with the clearcut only treatment having the slowest rate of basal area development.

### Stand Volume

Model [3] also was used to test responses of stand volume to treatment:

$$\ln(V) = 6.28 + 24.4/AGE - 1.68T1 - 1.09T2 - 0.326T3 - 0.617T4 \quad (6)$$

Stand volume differed significantly among site-preparation intensities, with the clearcut-only treatment having the




Figure 1—Effects of increasing site-preparation intensity or development of stand volume of loblolly pine.

of stand volume did not differ significantly ( $P \geq 0.05$ ) between treatments 5 and 6—the two treatments having the greatest rate of stand development.

## CONCLUSIONS

Effects of site preparation on stand development were modeled as adjustments to a growth model for stand average height, basal area, and volume. This approach provided an accurate description of stand responses to site-preparation intensity. Results suggest that the more intensive site preparation treatments lead to greater rates of stand development.

## LITERATURE CITED

- Clutter, Jerome L.; Fortson, James C.; Pienaar, Leon V. [and others]. 1983. Timber management: a quantitative approach. Malabar, FL: Krieger Publishing Company. 333 p.
- Glover, Glenn R.; Zutter, Bruce R. 1993. Loblolly pine and mixed hardwood stand dynamics for 27 years following chemical, mechanical, and manual site preparation. *Canadian Journal of Forest Research*. 23: 2126-2132.
- Harrington, Timothy B.; Edwards, M. Boyd. 1996. Structure of mixed pine and hardwood stands 12 years after various methods and intensities of site preparation in the Georgia Piedmont. *Canadian Journal of Forest Research*. 26: 1490-1500.
- Miller, James H.; Edwards, M. Boyd. 1985. Impacts of various intensities of site preparation on Piedmont soils after 2 years. In: Shoulders, Eugene, ed. Proceedings of the third biennial southern silvicultural research conference; 1984 November 7-8; Atlanta, GA. Gen. Tech. Rep. SO-54. Asheville, NC: U.S. Department of Agriculture, Forest Service, Southeastern Forest

level growth and yield response to environmental  
treatments. *Forest Science*. 41: 629-638.

closure of white spruce and lodgepole pine. *Canadian  
Journal of Forest Research*. 19: 262-269.

# **Proceedings of the Ninth Biennial Southern Silvicultural Research Conference**

***Edited by***

Thomas A. Waldrop

Clemson, South Carolina

February 25-27, 1997

***Hosted by***

Clemson University

USDA Forest Service, Southern Research Station

***Sponsored by***

Alabama A&M University

Clemson University

Historically Black Colleges and Universities Program

National Association of Consulting Foresters of America

National Association of Professional Forestry Schools and Colleges

National Hardwood Lumber Association

Society of American Foresters

Southern Group of State Foresters

Southern Industrial Forest Research Council

USDA Forest Service, Southern Research Station

University of Arkansas, Monticello

***Published by***

USDA Forest Service

Southern Research Station

Asheville, North Carolina